Note to Readers: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Suicide and Ambient Temperature in East Asian Countries: A Time-Stratified Case-Crossover Analysis

Yoonhee Kim, Ho Kim, Yasushi Honda, Yue Leon Guo, Bing-Yu Chen, Jong-Min Woo, and Kristie L. Ebi

Table of Contents

- **Table S1.** City-specific characteristics of suicide cases and population size.
- **Table S2.** Weather characteristics by season in fifteen cities.
- **Table S3.** The associations for temperature or sunshine duration.
- **Table S4.** City-specific associations between same-day temperature and suicide according to gender and age group.
- **Figure S1.** Yearly trend of suicide rates on national levels by gender in Korea, Japan, and Taiwan. Suicide and population data were obtained from Statistics Korea, Ministry of Strategy and Finance in South Korea, Ministry of Health, Labor and Welfare (for suicide) and Statistics Bureau, Ministry of Internal Affairs and Communications (for population) in Japan, and Department of Statistics, Ministry of Health and Welfare (for suicide) and Department of Statistics, Ministry of the Interior (for population) in Taiwan.
- **Figure S2.** Yearly trend of suicide rates on national levels by age groups in Korea, Japan, and Taiwan. Suicide and population data were obtained from Statistics Korea, Ministry of Strategy and Finance in South Korea, Ministry of Health, Labor and Welfare (for suicide) and Statistics Bureau, Ministry of Internal Affairs and Communications (for population) in Japan, and

Environ Health Perspect DOI: 10.1289/ehp.1409392

Department of Statistics, Ministry of Health and Welfare (for suicide) and Department of Statistics, Ministry of the Interior (for population) in Taiwan.

Figure S3. Monthly trend of suicide in fifteen cities. Study period varied depending on the country: Korea (1992–2010), Japan (1972–2010), and Taiwan (1994–2007). Suicide data were obtained from Statistics Korea, Ministry of Strategy and Finance in South Korea, Ministry of Health, Labor and Welfare in Japan, and Department of Statistics, Ministry of Health and Welfare in Taiwan.

Figure S4. Trends for day of the week of suicide in fifteen cities. Study period varied depending on the country: Korea (1992–2010), Japan (1972–2010), and Taiwan (1994–2007). Suicide data were obtained from Statistics Korea, Ministry of Strategy and Finance in South Korea, Ministry of Health, Labor and Welfare in Japan, and Department of Statistics, Ministry of Health and Welfare in Taiwan.

Figure S5. Lagged effect estimates for sunshine on suicide according to single lags. PC indicates percent change of suicide risk corresponding to a SD/2-increase of mean sunshine, adjusting for temperature, relative humidity, atmospheric pressure, long-term time-trend, and month.

Figure S6. Lagged effect estimates for sunshine on suicide according to moving averages. Moving average 90 indicates the average of sunshine for a week on previous ninety days. PC indicates percent change of suicide risk corresponding to a SD/2-increase of mean sunshine, adjusting for temperature, relative humidity, atmospheric pressure, long-term time-trend, and month.

Figure S7. Lagged effect estimates for temperature on suicide according to single lags. PC indicates percent change of suicide risk corresponding to a SD/2-increase of mean temperature, adjusting for sunshine duration, relative humidity, atmospheric pressure, long-term time-trend, and month.

Figure S8. Lagged effect estimates for temperature on suicide according to moving averages. PC indicates percent change of suicide risk corresponding to a SD/2-increase of mean temperature, adjusting for sunshine duration, relative humidity, atmospheric pressure, long-term time-trend, and month.